This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

◀



sFcRIIIA-G2 Variant 25°C Stability

B.

FcRillA-avi-blotin binding to IC:




FIG.




FIG. 2




FIG. 3




FIG. 4







FIG. 5










FIG. 6

B.




FIG. 7

Pharmacokinetics of sFcRIIB molecules in Balb/C mice




FIG. 8




FIG. 9






FIG. 10





- → · N treatment (N=6)

- → sFcRIIB-G2 N297Q 0.3ug/g (N=3)

FIG. 12





FIG. 13



Ô

10

20

-- N Treatment (N=3)

--- sFcRIIB-G2 N297Q 0.4ug/g (N=2)

→ 5.1 N297Q 0.25ug/g (N=2)

--- 5.1 N297Q 0.125ug/g (N=2)

- ■ - 22.1 N297Q 0.25ug/g (N=2)

- ▲ - 22.1N297Q 0.125ug/g (N=2)


В.

sFcRlIB-G2 N297Q and 5.1 N297Q

30

Hours

40



50

60

C.

sFcRIIB-G2 N297Q and 22.1 N297Q





FIG. 14



В.




FIG. 15




FIG. 16